

Stomp® Aqua

A herbicide for the control of a range of grass and annual broad-leaved weeds in a wide range of crops.

A capsule suspension containing 455 g/litre pendimethalin

Risk and Safety Information

Warning

**May cause an allergic skin reaction.
Suspected of damaging the unborn child.
Toxic to aquatic life with long lasting effects.**

If medical advice is needed, have product container or label at hand.
Avoid breathing mist/vapours/spray.
Wear protective gloves, protective clothing and eye protection or face protection.
Collect spillage.
Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. : 03750



® = Registered trademark of BASF

UFI : 66X2-POSP-P00S-AD7G

10 L

Supplied by:
BASF Ireland DAC, 4th and 5th
Floors, 2 Stockport Exchange,
Railway Road, Stockport, SK1 3GG,
UK Telephone: 00 44 (0)161 475 3000

Emergency Information
(24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0044 (0)845 602 2553
(office hours)

Authorisation holder:
BASF plc, 4th
and 5th Floors,
2 Stockport Exchange,
Railway Road,
Stockport, SK1 3GG,
UK Telephone:
00 44 (0)161 475 3000

81177339 IE 2084



IMPORTANT INFORMATION**FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE, as directed below:**

| Crops | Maximum Individual Dose | Maximum Number of Treatments | Latest Time of Application |
|--|--------------------------------|-------------------------------------|---|
| Winter wheat, Durum wheat, Winter barley, Winter rye and Triticale | 2.9 litres product/ha | One per crop | Before leaf sheath erect stage (crop GS 30). |
| Spring barley, potato, combining pea and sunflower | 2.9 litres product/ha | One per crop | Pre-crop emergence |
| Forage maize, grain maize | 3.3 litres product/ha | One per crop | Before 4th leaf stage |
| Bulb onion (spring and autumn, drilled and transplanted). | 2.9 litres product/ha | One per crop | Pre-crop emergence |
| Leek | 2.9 litres product/ha | One per crop | Pre-crop emergence |
| Carrot and parsnip | 2.9 litres product/ha | One per crop | Pre-crop emergence |
| Broccoli/calabrese, Brussels sprout, Cabbage, Cauliflower | 2.9 litres product/ha | One per crop | Before transplanting |
| Blackcurrant, Gooseberry | 2.9 litres product/ha | One per year | Before bud burst |
| Strawberry (outdoor) | 2.9 litres product/ha | One per year | After flower initiation but before flower truss emergence |
| Apple, Cherry, Pear, Plum | 2.9 litres product/ha | One per year | Before bud burst |
| Raspberry, Loganberry, Rubus Hybrid, Blackberry (outdoor) | 2.9 litres product/ha | One per year | After harvest but before bud burst |

READ ALL PRECAUTIONS BEFORE USE.**PRECAUTIONS****Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS) AND SUITABLE PROTECTIVE GLOVES when applying by hand held equipment.

WASH ALL PROTECTIVE CLOTHING thoroughly after use, especially the insides of gloves.

WASH CONCENTRATE from skin or eyes immediately.

WASH HANDS AND EXPOSED SKIN before meals and after work.

AVOID ALL CONTACT BY MOUTH.

DO NOT BREATHE SPRAY.

Environmental protection

Extreme care should be taken to avoid damage by drift onto plants outside the target area.

Do not contaminate water with the product or its container. (Do not clean application equipment near surface water/Avoid contamination via drains from farmyards and roads).

To protect aquatic organisms respect an unsprayed buffer zone of 5m to surface water bodies.

Do not allow spray from handheld sprayers to fall within 1m of the top of the bank of a static or flowing waterbody.

Aim spray away from water.

STORAGE

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times.

Keep dry and frostproof in a suitable pesticide store.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

A herbicide for the control of annual-grass and broad-leaved weeds in a wide range of crops.

Restrictions/Warnings

Efficacy

Some soil moisture must be present for STOMP AQUA to be activated. Best results will be obtained if rainfall occurs within seven days of application.

Residual control may be reduced:

- under prolonged dry conditions
- on soils with a high Kd factor
- where organic matter exceeds 6%
- where ash content is high.

Do not disturb the soil after STOMP AQUA has been applied as this will result in reduced weed control. Where cultural techniques which encourage the build up of organic residues in the surface soil are practised for a number of seasons, the effectiveness of residual herbicides may be reduced. In such circumstances periodic ploughing is recommended to disperse residues into a greater volume of soil.

Soil types

STOMP AQUA may be used on all mineral soil types.

Do not use on soils with more than 10% organic matter.

On stony or gravelly soils there is a risk of crop damage, especially if heavy rain falls soon after application.

Do not use on water logged soil or soils prone to water logging.

Seedbed preparation

Trash and straw should be incorporated evenly during seedbed preparation.

Seedbed must have a fine, firm tilth.

Consolidate loose or cloddy seedbeds before use.

Following pre-emergence applications, unconsolidated clods (especially if larger than 75 mm (3" diameter) may reduce the level of weed control and cause seed to be inadequately covered, which could result in crop damage.

Crop safety

Extreme care should be taken to avoid damage by drift onto plants outside the target area.

Do not apply STOMP AQUA to crops suffering from stress, which may be caused for example by pests, disease, water logging, poor seedbed conditions or previous chemical treatment.

Seed should be covered with a minimum of 3.2 cm of settled soil (2.5 cm for Peas and Sunflowers, 5 cm for Maize).

Shallow drilled crops should be treated post-emergence.

Do not soil incorporate.

Do not spray undersown crops.

Do not undersow crops treated with STOMP AQUA.

STOMP AQUA should not be used on protected crops, or in greenhouses.

Other Restrictions/Warnings

For professional use only.

Before using STOMP AQUA on crops to be processed please consult your processor.

Concentrated or diluted STOMP AQUA will stain. Avoid spillage.

Staining is minimised or completely removed if skin and clothes are washed immediately.

Hose down machinery immediately after use with a spray tank cleaner.

Weed control

Cereals

STOMP AQUA applied alone

All weed susceptibility ratings in the table below are for applications made pre-emergence of the weeds.

| CROPS | Winter Wheat, Durum Wheat, Winter Barley, Winter Rye & Triticale | | Spring Barley |
|---------------------------|--|-----|---------------|
| PRODUCT | STOMP AQUA | | STOMP AQUA |
| RATE (litres/ha) | 2.9 | 2.2 | 2.9 |
| GRASS WEED CONTROL | | | |
| Annual Meadow-grass | S | S | S |
| Awned Canary Grass | - | - | - |
| Black-grass | - | - | - |
| Rough Meadow-grass | MS | MS | MS |
| BROAD-LEAVED WEEDS | | | |
| Black-bindweed | - | - | - |
| Black Nightshade | - | - | - |
| Cleavers | - | - | - |
| Common Chickweed | S | S | S |
| Common Fumitory | MS | - | MS |

| CROPS | Winter Wheat, Durum Wheat, Winter Barley, Winter Rye & Triticale | | Spring Barley |
|------------------------------|--|-----|---------------|
| PRODUCT | STOMP AQUA | | STOMP AQUA |
| RATE (litres/ha) | 2.9 | 2.2 | 2.9 |
| Common Orache | S | MS | S |
| Common Poppy | S | MS | S |
| Corn Buttercup | - | - | - |
| Corn Marigold | S | S | S |
| Fat-hen | S | MS | S |
| Field Forget-me-not | S | MS | S |
| Field Pansy | S | MS | S |
| Hemp-nettle (Day Nettle) | S | S | S |
| Henbit Dead-nettle | S | S | S |
| Knotgrass | S | MS | S |
| Mayweeds | MS | - | - |
| Parsley Piert | S | S | S |
| Red Dead-nettle | S | S | S |
| Redshank (early germinating) | - | - | - |
| Scarlet Pimpernel | S | S | S |
| Shepherd's Purse | MS | - | MS |
| Small Nettle | S | - | S |
| Smooth Sowthistle | S | MS | S |
| Speedwells | S | S | S |
| Volunteer Oilseed Rape (1) | S | S | MS |

S = Susceptible

MS = Moderately susceptible

(1) = Deep germinating Volunteer Oilseed Rape may not be controlled

- = no data

Other crops

All weed susceptibility ratings in the table below are for applications made pre-emergence of the weeds.

| CROPS | Combining peas, Sunflowers, Carrots, Parsnips, Strawberries, Bush fruit, Cane fruit, Top fruit | Onions, Leeks, Transplanted Brassicas: – (Brussels sprouts, Broccoli, Cabbages, Calabrese, Cauliflowers | Forage Maize, Grain Maize | Potatoes : –(First Early, Second Early, Maincrop) |
|--|--|---|---------------------------|--|
| PRODUCT RATE (litres/ha) Tank mix partner | STOMP AQUA 2.9 | STOMP AQUA 2.9 | STOMP AQUA 3.3 | STOMP AQUA 2.9 Approved metribuzin product to give 350 g/ha metribuzin |
| Annual Meadow-grass | S | S | S | S |
| Black-grass | – | – | – | – |
| Rough Meadow-grass | MS | MS | MS | MS |
| Black & White Mustard | – | – | – | – |
| Black bindweed | – | – | – | MS |
| Black Nightshade | – | – | S # | – |
| Charlock | – | – | – | S |
| Cleavers (#) | – | – | – | MS |
| Common Chickweed | S | S | S | S |
| Common Fumitory (#) | *MS | MS | MS | MS |
| Common Orache | S | S | S | S |
| Common Poppy | S | S | S | S |
| Corn Buttercup | – | – | – | – |
| Corn Marigold | S | S | S | S |
| Fat-hen | S | S | S | S |
| Field Forget-me-not | S | S | S | S |
| Field Pansy | S | S | S | S |
| Groundsel | – | – | – | S |
| Hemp (Day) -nettle | S | S | S | S |
| Henbit Dead-nettle | S | S | S | S |
| Knotgrass | S | S | S | S |
| Mayweeds (#) | – | – | – | S |
| Parsley Piert | S | S | S | S |
| Red Dead Nettle | S | S | S | S |
| Redshank (1) | – | – | – | S |
| Scarlet Pimpernel | S | S | S | S |

| | | | | |
|--|--|---|---------------------------|--|
| CROPS | Combining peas, Sunflowers, Carrots, Parsnips, Strawberries, Bush fruit, Cane fruit, Top fruit | Onions, Leeks, Transplanted Brassicas: – (Brussels sprouts, Broccoli, Cabbages, Calabrese, Cauliflowers | Forage Maize, Grain Maize | Potatoes : –(First Early, Second Early, Maincrop) |
| PRODUCT RATE (litres/ha) Tank mix partner | STOMP AQUA 2.9 | STOMP AQUA 2.9 | STOMP AQUA 3.3 | STOMP AQUA 2.9 Approved metribuzin product to give 350 g/ha metribuzin |
| Shepherd's Purse | *MS | MS | MS | MS |
| Small Nettle | S | S | S | S |
| Smooth Sowthistle | S | S | S | MS |
| Speedwells | S | S | S | S |
| Volunteer Oilseed Rape (2) | MS | MS | MS | S |

S = Susceptible

MS = Moderately susceptible

* = Control may be achieved under favourable conditions

(1) = Early germinating

(2) = Deep germinating Volunteer Oilseed Rape may not be controlled

– = no data

= If application is followed by a period of dry conditions, or in situations where very heavy populations occur, a sequence of STOMP AQUA and a product applied post-emergence may be necessary.

Resistance Management

GROUP

3

HERBICIDE

Strains of some annual grasses (eg Black-grass, Wild-Oats, and Italian Ryegrass) have developed resistance to herbicides, which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from Teagasc, your distributor, crop adviser or product manufacturer. Populations of Black-grass and Italian ryegrass with high levels of enhanced metabolism resistance will not be fully controlled.

Key elements of the resistance management strategy for STOMP AQUA:

- Always follow WRAG guidelines for preventing and managing herbicide resistant weeds.
- Maximise the use of cultural control measures wherever possible (e.g. crop rotation, ploughing, stale seedbeds, delayed drilling, etc).
- Use tank mixes or sequences of effective herbicides with different modes of action within individual crops, or successive crops.
- For the control of herbicide resistant grassweeds, always use STOMP AQUA in tank mix or sequence with other effective graminicides with different modes of action.

- Apply pre-emergence of weeds wherever possible. If applications are delayed, apply post-emergence products/mixtures to small, actively growing weeds, especially where high levels of resistance are suspected and to reduce the risk of resistance development.
- Monitor fields regularly and investigate the reasons for any poor control.

Crop Specific Information

STOMP AQUA is recommended for use on all varieties of approved crops on any mineral soil except where indicated in the table below.

Winter wheat including Durum wheat, winter barley

| | |
|--------------|---|
| Product | STOMP AQUA |
| Rate | 2.9 or 2.2 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Pre-emergence of the crop to before leaf sheath erect stage, (GS30). Do not apply pre-emergence to crops drilled after 30th November. |
| Seed depth | Seed must be covered with a MINIMUM of 3.2cm of settled soil. ONLY treat shallow drilled crops POST-EMERGENCE. |

Winter rye, Triticale

| | |
|--------------|---|
| Product | STOMP AQUA |
| Rate | 2.9 or 2.2 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Pre-emergence of the crop to before leaf sheath erect stage, (GS30). Do not apply pre-emergence to crops drilled after 30th November. |
| Seed depth | Seed must be covered with a MINIMUM of 3.2cm of settled soil. ONLY treat shallow drilled crops POST-EMERGENCE. |

Spring barley

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Pre-emergence of the crop. Apply STOMP AQUA as soon as possible after drilling and before emergence. Due to risk of dry soils, do not apply STOMP AQUA alone after the end of March unless rainfall is imminent. |
| Seed depth | Seed must be covered with a MINIMUM of 3.2 cm of settled soil. |

Combining Peas

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Pre-emergence of the crop. Apply STOMP AQUA as soon as possible after sowing and final seedbed preparation. Do not apply if the plumule is less than 13mm from the soil surface. Due to risk of dry soils, do not apply STOMP AQUA alone after the end of March unless rainfall is imminent. |
| Soil types | All mineral soils except gravelly soil |
| Seed depth | Seed must be covered with a MINIMUM of 2.5cm of settled soil. |

Potatoes (First early, second early & maincrops)

| | |
|--------------|--|
| Products | STOMP AQUA + Approved metribuzin product to give 350 g/ha metribuzin. |
| Rate | 2.9 l/ha + appropriate rate to deliver 350g/ha metribuzin In dry conditions apply a STOMP AQUA – metribuzin sequence. |
| Water volume | 200 litres/hectare. |
| Timing | Pre-emergence of the crop. Apply as soon as possible after planting and final ridging up. Loose structured ridges must be allowed time for settlement before application. Do not apply later than 7 days before emergence. |
| Soil types | Do not use on Sands (S), Gravelly or Stony soils. |
| Variety | Read the Metribuzin product label carefully, particularly with regard to varietal restrictions. |
| Application | STOMP AQUA should be applied in a minimum of 200 litres of water/ha. |
| Notes | Best weed control will be achieved with settled well-rounded ridges with few clods. If re-ridging is necessary, delay application until after the final ridging is completed. Slight distortion and discolouration of the initial shoots may occur if very heavy rain falls after application but before emergence, particularly to crops grown on very light soils. This is quickly outgrown and subsequent growth is unaffected. Read the Metribuzin product label carefully, particularly with regard to following crop restrictions. |

Sunflowers

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply as soon as possible after sowing and final seedbed cultivation, before crop and weed emergence. |
| Seedbed | Consolidate seedbeds after drilling to provide a firm level soil. Seed should be drilled so that after seedbed consolidation it is covered by a minimum of 2.5 cm of settled soil. |

Carrots and Parsnips

| | |
|--------------|---|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply as soon as possible after drilling but before crop and weed emergence. |
| Notes | If emerged weeds are present after drilling but pre-emergence of the crop, STOMP AQUA may be applied in tank mix with a recommended approved contact herbicide. |

Transplanted Brassicas (Broccoli, Brussels sprouts, Cabbages, Calabrese, Cauliflowers)

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply after final plantbed cultivation but before transplanting. Do not apply STOMP AQUA post-planting as crop damage may occur. Do not apply STOMP AQUA to any transplanted brassicas when heavy rain is forecast. |
| Application | Do not incorporate and avoid all unnecessary disturbance to soil after application. When transplanting care must be taken not to introduce treated soil into the root zone. If necessary, irrigation should be used before application as some moisture is essential for the chemical to be activated. |
| Soil types | Do not use on crops grown on sands (CS, S, FS, LCS), very light soils (LS, LFS, CSL), as crop damage may result. |
| Notes | If emerged weeds are present at pre-transplanting application, apply STOMP AQUA in tank mix with a recommended approved contact herbicide. |

Bulb onions (Spring and autumn drilled or transplanted) and Leeks (drilled only)

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Drilled crops – apply STOMP AQUA as soon as possible after drilling but before crop or weed emergence. Transplanted crops – apply STOMP AQUA pre-transplanting. Do not apply STOMP AQUA to any onion or leek crop when heavy rain is forecast. |
| Seed depth | Seed should be covered with a minimum of 2.5cm of settled soil. |
| Application | When transplanting, care must be taken not to introduce treated soil into the root zone. |
| Soil types | STOMP AQUA is not recommended for use on onions or leeks grown on sands (CS, S, FS, LCS), very light soils (LS, LFS, CSL), as crop damage may result. STOMP AQUA is not recommended for use on onions or leeks grown on fen soils or other soils containing in excess of 10% organic matter, as weed control may be reduced. |
| Notes | If weeds are present, these can be controlled by applying STOMP AQUA in tank mix with a recommended contact herbicide. Read tank mix partner label carefully for restrictions on transplanted multi-seeded onions or leeks. |

Forage maize, Grain maize

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 3.3 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Pre-emergence to before 4th leaf of the crop. |
| Notes | Do not use on Sweetcorn or Maize grown for seed. Seed must be covered by a minimum of 5 cm of settled soil. The use of STOMP AQUA may affect the full development of crown roots which function only to anchor the plant. This has no effect on the yield of maize. If application is followed by a period of dry conditions or in situations where very heavy weed populations occur, a sequence of STOMP AQUA and a product applied post-emergence may be necessary. |

Strawberries (outdoor, maiden and established crops)

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | <p>Maiden Beds</p> <ul style="list-style-type: none"> runners should be planted so that roots are well covered. good consolidation of the plantbed is necessary for good weed control. STOMP AQUA should be applied to dormant newly planted runners in the autumn or early spring immediately after planting and prior to weed emergence. if runners are likely to be slow in establishing due to stress conditions, such as drought, at the time of planting, the application of STOMP AQUA should be delayed until plants have established and are free of stress. application made after runners have started growing away may reduce the initial vigour of new foliage but this will be rapidly outgrown. do not apply STOMP AQUA before October to beds newly planted with cold-stored runner or beds newly planted in late summer. pre-planting application of STOMP AQUA is not recommended. <p>Established beds</p> <ul style="list-style-type: none"> apply STOMP AQUA to weed free soil from autumn to early spring during the dormant period of the crop. STOMP AQUA applied after the end of March or after the emergence of flower trusses in the spring may affect crop yield, particularly if conditions adverse to vigorous plant growth follow application. do not apply STOMP AQUA during the flower initiation period (immediately post-harvest to mid-September). |
| Notes | Leaf growth of strawberries may be checked following applications of STOMP AQUA in the spring but, in extensive experimentation, this has been shown not to affect yield. Do NOT use STOMP AQUA on protected crops or crops grown in green houses. |

Bush Fruit – Blackcurrants, Gooseberries

| | |
|--------------|---|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply STOMP AQUA after final cultivation to weed-free soil from autumn to early spring during the dormant period of the crop before bud burst, either over the top or as a directed spray. STOMP AQUA is not recommended for use in the season of planting. |
| Tank mixes | If emerged weeds are present at application, STOMP AQUA may be applied as a directed spray in tank mix with a recommended approved contact herbicide. |

Cane Fruit – Raspberries, Loganberries, Tayberries, Blackberries

| | |
|--------------|--|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply STOMP AQUA after final cultivation to weed-free soil from autumn to early spring. STOMP AQUA is not recommended for use after the end of March unless adequate soil moisture is present as some soil moisture present is essential for the chemical to be activated. Do not apply to autumn fruiting raspberries. Newly planted crops: STOMP AQUA should be applied immediately after planting. A light ridging along the cane row before application is recommended to ensure roots are well covered. STOMP AQUA should be applied well before the emergence of the new canes. Established crops: STOMP AQUA should be applied as soon as the canes have been cut out and tied, but before bud burst |
| Tank mixes | If emerged weeds are present at application, STOMP AQUA may be applied as a directed spray in tank mix with a recommended approved contact herbicide. |

Top Fruit – Apples, Cherries, Pears, Plums

| | |
|--------------|---|
| Product | STOMP AQUA |
| Rate | 2.9 l/ha |
| Water volume | 100 to 200 litres/hectare. |
| Timing | Apply STOMP AQUA to weed-free soil from autumn to early spring. STOMP AQUA is not recommended for use after the end of March unless adequate soil moisture is present as some soil moisture is essential for the chemical to be activated. STOMP AQUA is not recommended for use in the season of planting. |
| Tank mixes | If emerged weeds are present at the time of application, STOMP AQUA should be applied in tank mix with a recommended approved contact herbicide. |

Following Crops

Following crops after normal harvest

Before Rye grass is drilled after a very dry season, plough or cultivate to at least 15 cm.

If spring crops are to be followed by crops other than cereals plough or cultivate to at least 15cm.

In the event of crop failure

In the event of crop failure the land must be ploughed or thoroughly cultivated to a minimum depth of 15cm to ensure any residues are evenly dispersed throughout the soil.

The minimum intervals (specified below) should elapse between application of STOMP AQUA and the sowing of one of the following crops listed below.

| Application timing | Minimum interval | In the event of crop failure, the following crops may be drilled: |
|-----------------------|------------------|--|
| Autumn | 5 months | Spring wheat, Spring barley, Spring Field beans, Broad beans, Dwarf beans, Brussels sprouts, Cabbage, Calabrese, Carrots, Cauliflower, Parsnips, Parsley, Peas, Potato, Linseed, Maize, Turnip |
| Spring & early summer | 2 months | Spring Field beans, Broad beans, Dwarf beans, Brussels sprouts, Cabbage, Calabrese, Carrots, Cauliflower, Parsnips, Parsley, Peas, Linseed, Turnip |
| | 5 months | Any crop (with the exception of Red Beet, Sugar Beet and Spinach) may be planted or sown. |
| | 12 months | Red Beet, Sugar Beet and Spinach |

Mixing and Application

Mixing

Never prepare more spray solution than is required.

Three quarters fill the tank with clean water and start the agitation. To ensure thorough mixing of the product, invert the container several times before opening. Add the required quantity of STOMP AQUA to the spray tank while re-circulating. Fill up the tank with water and continue agitation until spraying is completed.

On emptying the container, rinse container thoroughly by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of container safely.

When tank mixes are to be used each product should be added separately to the spray tank and fully dispersed before the addition of any further product(s). Add products to the spray tank in the following order ;

1. wettable powders (WP)
 2. dry flowables or water dispersible granules (WG)
 3. suspensions (e.g. STOMP AQUA)
 4. suspension concentrates (e.g. SC or SE)
 5. emulsifiable concentrates (e.g. EW or EC)
 6. soluble concentrates (SL)
- Maintain agitation at all times.
 - Spray out as soon as possible after mixing.
 - Do not let the mixture stand.

Application

Ensure good, even spray cover of the target using a FINE or MEDIUM quality spray, as defined by BCPC.

Do not use less than 50 mesh inline boom or nozzle filters.

For boom sprayers

Apply STOMP AQUA in 100-200 l/ha. For potatoes apply STOMP AQUA in minimum 200 l/ha water.

When tank mixing with other products use a minimum water volume of 150-200 l/ha depending on the tank mix partner.

For knapsack sprayers

Use a maximum of 15 mls of STOMP AQUA per litre of water and ensure a good, fine coverage of the target.

Sprayer cleaning

After spraying, thoroughly clean and flush out application machinery with a minimum of three rinses, to ensure that all traces of product are removed to avoid sticking of dried material to spray tank walls or spray lines, etc.

Tank Mixtures

Please contact your agronomist, supplier or the BASF Technical Hotline (0044 (0)845 602 2553) for up to date information on the compatibility of Stomp Aqua.

Sequential Treatments

STOMP AQUA may be used in sequence with any other approved product. Leave a minimum interval of 24 hours unless longer is specified on the label.

STOMP AQUA may be applied in sequence with Avadex BW granular provided only one product is applied pre-emergence of the crop.

Trade Mark Acknowledgments

STOMP is a registered trademark of BASF.

Avadex is a registered trademark of Gowan Internacional Limitada.

The following does not form part of the product label under S.I. No. 159 of 2012

With many products there is a general risk of resistance developing to the active ingredients. For this reason a change in activity cannot be ruled out. It is generally impossible to predict with certainty how resistance may develop because there are so many crop and use connected ways of influencing this. We therefore have to exclude liability for damage or loss attributable to any such resistance that may develop. To help minimise any loss in activity the BASF recommended rate should in all events be adhered to.

Numerous, particularly regional or regionally attributable, factors can influence the activity of the product. Examples include weather and soil conditions, crop plant varieties, crop rotation, treatment times, application amounts, admixture with other products, appearance of organisms resistant to active ingredients and spraying techniques. Under particular conditions a change in activity or damage to plants cannot be ruled out. The manufacturer or supplier is therefore unable to accept any liability in such circumstances. All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their mixing or use or the weather conditions during and after application, which may affect the performance of the material, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use; but nothing should be deemed to exclude or restrict any liability upon us which cannot be excluded or restricted under the provisions of any applicable law.

Additional Product Safety Information

The information on this label is based on the best available information including data from test results.

Safety Data Sheet

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



agricentre.basf.ie/Stomp-Aqua-IE/MSDS

Alternatively, contact your supplier.

Stomp® Aqua

A herbicide for the control of a range of grass and annual broad-leaved weeds in a wide range of crops.

A capsule suspension containing 455 g/litre pendimethalin

Risk and Safety Information

Warning

**May cause an allergic skin reaction.
Suspected of damaging the unborn child.
Toxic to aquatic life with long lasting effects.**

If medical advice is needed, have product container or label at hand.
Avoid breathing mist/vapours/spray.
Wear protective gloves, protective clothing and eye protection or face protection.
Collect spillage.
Store locked up.

Dispose of contents/container to a licensed hazardous waste disposal contractor or collection site except for triple rinsed empty containers which can be disposed of as non-hazardous waste.

To avoid risks to human health and the environment, comply with the instructions for use.

PCS No. : 03750



UFI : 66X2-P0SP-P00S-AD7G

10 L

Supplied by:
BASF Ireland DAC, 4th and 5th
Floors, 2 Stockport Exchange,
Railway Road, Stockport, SK1 3GG,
UK Telephone: 00 44 (0)161 475 3000
Emergency Information
(24 hours freephone): 0049 180 227 3112
Technical Enquiries: 0044 (0)845 602 2553
(office hours)

Authorisation holder:
BASF plc, 4th
and 5th Floors,
2 Stockport Exchange,
Railway Road,
Stockport, SK1 3GG,
UK Telephone:
00 44 (0)161 475 3000