

Document size: 114x67 mm (width x height)
Gross size: 114x67 mm (width x height)

Storm[®] Ultra Secure



BASF

We create chemistry

Advanced, highly attractive block bait.
Excellent and fast control in all situations.

275 g



® = Registered Trademark of BASF

Block Bait – Controls rats and mice

- ✓ Use in a tamper-resistant bait station
- ✓ For use indoor and outdoor around buildings
- ✓ Securable



81178618 GB 2104

Storm® Ultra Secure

Contains 0.025g/kg (0.0025% w/w) flocoumafen.

Storm Ultra Secure is a ready-to-use block bait containing flocoumafen (0.0025% w/w) for use as a rodenticide by the General Public for the control of rats and mice indoors and outdoors around buildings for the protection of public health, stored products and materials.

Storm Ultra Secure is for use in and around buildings only.

For general public use.

This product contains a bittering agent and a dye.

Great Britain Authorisation Holder:

BASF plc,
4th and 5th Floors, 2 Stockport Exchange,
Railway Road, Stockport, SK1 3GG, UK

Telephone: 0161 475 3000

Emergency information: 0049 180 2273112 (24 Hours freephone)

GB Authorisation number:

GB-2018-1164-0001

® = Registered trademark of BASF

Ireland and Northern Ireland Authorisation Holder:

BASF Ireland DAC

Asgard House, 19-20 City Quay

D02 K744, Dublin, Ireland

Telephone: 0044 (0) 161 475 3000

Emergency information: 0049 180 2273112 (24 Hours freephone)

Ireland Authorisation number:

IE/BPA 70694-001

Northern Ireland Authorisation number:

NI-2018-1164-0001

Expiry date:

3 years after date of manufacture.

Batch number:

see packaging.



First aid:

This product contains an anticoagulant substance. If ingested, symptoms, which may be delayed, may include nosebleed and bleeding gums. In severe cases, there may be bruising and blood present in the faeces or urine.

Antidote: Vitamin K1 administered by medical/veterinary personnel only.

In case of:

- Dermal exposure, wash skin with water and then with water and soap.
- Eye exposure, rinse eyes with eyes-rinse liquid or water, keep eye lids open at least 10 minutes.
- Oral exposure, rinse mouth carefully with water. Never give anything by mouth to unconscious person. Do not provoke vomiting. If swallowed, seek medical advice immediately and show the product's container or label.

Poison Information

UK: UK medical professionals should contact the National Poisons Information Service (www.npis.org) for further advice.

IE: For information or to report a poisoning incident contact The National Poisons Information Centre, Beaumont Hospital, Dublin (01-809 2166), retain the label for reference.

Contact a veterinary surgeon in case of ingestion by a pet.

To access the Safety Data Sheet for this product scan the QR code or use the weblink below:



UK:

<https://pest-control.basf.com/gb/en/products/rodenticides/storm-ultra-secure#downloads>



IE:

agricentre.basf.ie/Storm-Ultra-Secure-IE/MSDS

Alternatively, contact your supplier.

Application method:

Place Storm Ultra Secure baits throughout the infested area using tamper-resistant bait stations only.

UK – Tamper-resistant bait stations must be:

- Strong enough to prevent entry or destruction by dogs.
- Strong enough to prevent entry or destruction by children under six years of age using hands, feet or objects reasonably expected to be available in the use area (e.g. household objects, toys, sticks, stones etc.).
- Lockable and sealable so that children and dogs cannot gain access through the opening or mechanisms used to fill the bait compartments.
- Resistant to destruction or weakening from exposure to typical non-catastrophic weather (e.g. direct sunlight, extremes of temperature and humidity, rain, snow etc.).

Application rate and frequency:

Species	Situation	Application rate for each bait station	Frequency
House mouse (<i>Mus musculus</i>)	Indoor	25 g (1 block) If more than one bait station is needed, the distance between bait stations should be 1 - 2 meters.	The bait stations should be visited at least every 2 to 3 days at the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary.

Species	Situation	Application rate for each bait station	Frequency
Brown rat (<i>Rattus norvegicus</i>)	Indoor and outdoor around buildings	50 g to 75 g (2-3 blocks).	The bait stations should be visited only 5 to 7 days after the beginning of the treatment and at least weekly afterwards, in order to check whether the bait is accepted, the bait stations are intact and to remove rodent bodies. Re-fill bait when necessary
Black or roof rat (<i>Rattus rattus</i>)		If more than one bait station is needed, the distance between bait stations should be 5 - 10 meters.	

Instructions for use:

Read and follow the product information as well as any information accompanying the product or provided at the point of sale before using it.

Prior to the use of rodenticide products, non-chemical control methods (e.g. traps) should be considered.

Remove food which is readily attainable for rodents (e.g. spilled grain or food waste). Apart from this, do not clean up the infested area just before the treatment, as this only disturbs the rodent population and makes bait acceptance more difficult to achieve.

The product should be placed in the immediate vicinity of places where rodent activity has previously occurred (e.g. rodent runs, nesting sites, holes, burrows etc.).

Do not place bait stations near water drainage systems or close to surface waters (e.g. rivers, ponds, water channels, dykes, irrigation ditches) where they can come into contact with water.

Protect bait from atmospheric conditions (e.g. rain, snow, etc.). Place the bait stations in areas not liable to flooding.

Where possible, bait stations must be fixed to the ground or other structures.

Place bait stations out of the reach of children, birds, pets, farm animals and other non-target animals.

Place bait stations away from food, drink and animal feeding stuffs, as well as from utensils or surfaces that have contact with these.

When using the product do not eat, drink or smoke. Wash hands and directly exposed skin after using the product.

Replace any bait in a bait station in which bait has been damaged by water or contaminated by dirt.

Remove the remaining bait or the bait stations at the end of the treatment period.

Because of their delayed mode of action, anticoagulant rodenticides take from 4 to 10 days to be effective after consumption of the bait.

UK - Risk mitigation measures:

Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Using this product should eliminate rodents within 35 days. In case of suspected lack of efficacy by the end of the treatment (i.e. rodent activity is still observed), the user should seek advice from the product supplier or call a pest control service.

Bait stations must be labelled with the following information: "do not move or open"; "contains a rodenticide"; "product name or authorisation number"; "active substance(s)" and "in case of incident, call a poison centre".

Hazardous to wildlife.

Search for and remove dead rodents during treatment, at least as often as bait stations are inspected.

Dispose of dead rodents in accordance with local requirements.

In the UK, poisoned rodents should be double-bagged using plastic bags and either disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or collected by a specialist waste contractor or the local authority.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

UK - Disposal:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with local requirements.

In the UK waste bait should be double bagged in plastic bags and disposed of in a household waste bin with a secure lid to prevent access of wildlife or pets or taken to a civic amenity site. For information on civic amenity sites contact the local authority.

IE - Risk mitigation measures:

Consider preventive control measures (plug holes, remove potential food and drinking as far as possible) to improve product intake and reduce the likelihood of reinvasion.

Do not use baits containing anticoagulant active substances as permanent baits for the prevention of rodent infestation or monitoring of rodent activities.

Where control of rodents has not been achieved by a control program after 35 days, the user should seek advice from the product supplier or call a pest control service.

Bait stations must be labelled with the following information: "Product name or authorisation number"; "Active substance(s)"; "Contains a rodenticide"; "Do not move or open" and "In case of incident, call the National Poisons Information Centre on (01) 809 2166".

Hazardous to wildlife.

Rodents can be disease carriers. Do not touch dead rodents with bare hands, use gloves or use tools such as tongs when disposing them.

IE - Disposal of uneaten bait and dead rodents:

At the end of the treatment, dispose of uneaten bait and the packaging in accordance with EPA requirements for the disposal of hazardous waste.

Use of gloves is recommended.

Dispose of dead rodents in accordance with local requirements. In Ireland, poisoned rodents should be double-bagged using plastic bags and either disposed of in a house-hold waste bin with a secure lid to

prevent access of wildlife or pets, or collected by a specialist waste contractor.

IE - Use Biocides Safely and Sustainably.

It is illegal to use this product for uses or in a manner other than prescribed on this label.

Storage:

Store in a dry, cool and well ventilated place.

Keep the container closed and away from direct sunlight.

Store in places prevented from the access of children, birds, pets and farm animals.

Shelf life: 3 years.